



Clinical Lab Intelligence

EDITION 007 | SEPTEMBER 7, 2026

From bench decisions to system design

This issue connects morphology, toxicology, microbiology stewardship, endocrine interpretation, MLS education, environmental quality systems, and emerging point-of-care engineering.

READING KEY Finding = what the source established. Why it matters = practice, education, or leadership relevance. Limits = what should not be overclaimed.	COVERAGE CHECK No DOI, canonical source, or substantially identical story from Editions 001-006 is repeated. This advance-prepared issue uses sources verified through September 2.
--	---

Core developments

01 AI assistance improved routine peripheral-blood morphology performance

Lu et al. | npj Digital Medicine | August 5, 2026 | Peer-reviewed; open access | [SOURCE](#)

Finding. In a prospective randomized paired study of 1,570 smears at three centers, AI-assisted review outperformed manual microscopy across 15 nucleated-cell categories. It improved abnormal-promyelocyte screening for acute promyelocytic leukemia (F1 0.794 vs 0.658), reduced platelet-estimate deviation from flow cytometry (2.05 vs 14.08 x 10⁹/L), and increased technician efficiency by about 60%.

Why it matters. This is operational evidence for assisted morphology, not merely an algorithm benchmark. Laboratories need validation plans that measure technologist correction burden, critical-cell sensitivity, failure escalation, platelet-estimate agreement, and time saved without weakening review competency.

Limits / cautions. One vendor system and three centers in China; the accepted version may receive editorial changes. Mindray funded the regulatory study, supplied equipment, and employed two authors. AI remained an aid, not an autonomous diagnosis.

02 The safer toxicology assay depended on the laboratory workflow

Jackson et al. | The Journal of Applied Laboratory Medicine | August 19, 2026 | Peer-reviewed | [SOURCE](#)

Finding. Across 238,705 urine EtG tests from separate implementation periods, the ARK assay produced more false positives (1.35% vs 0.15%), while the Immunalysis assay produced more interference flags (2.86% vs 0.45%). The laboratory selected the higher-interference method because flagged and presumptive-positive results could be routed to LC-MS/MS confirmation.

Why it matters. Method selection is a systems decision. Supervisors should compare analytical behavior together with reflex capacity, reporting language, turnaround time, confirmatory cost, client interpretation, and the potential consequences of an incorrect alcohol-use result.

Limits / cautions. The assays were not run concurrently on the same patients; negative screens were not mass-spectrometry confirmed; patient histories and interferent identities were unavailable.

03 Selective respiratory-panel use cut testing by more than 95%

Jenkins et al. | Microbiology Spectrum | July 28, 2026 | Peer-reviewed; open access | [SOURCE](#)

Finding. After a safety-net system removed unrestricted ordering and adopted targeted approval criteria, median multiplex respiratory-panel volume fell from 83 to 3 tests per week, a 96.4% decline sustained for one year. Annualized laboratory cost fell from \$741,446 to \$30,623 while overall positivity remained similar.

Why it matters. The intervention combined menu governance, EHR design, clinician communication, and selective access. Laboratory leaders can use this as a model for measuring low-value testing, designing exceptions, and monitoring unintended substitution after de-implementation.

Limits / cautions. Single-system retrospective quasi-experiment. It demonstrates utilization and cost change, not equivalent patient outcomes, antibiotic use, missed diagnoses, or transportability to every population.

04 Microbiology leaders see IT and clinical partnerships as the stewardship engine

She et al. | Journal of Clinical Microbiology | June 22, 2026; August issue | Peer-reviewed; open access | [SOURCE](#)

Finding. Among 119 clinical microbiology respondents, 73.9% reported practicing diagnostic stewardship some or most of the time. IT was rated the most effective strategy by 78.2%, while time and personnel were the leading barrier (52.1%); academic centers more often had formal consultation, utilization committees, or formularies.

Why it matters. Stewardship cannot rely on individual memory. A practical leadership agenda includes order-entry rules, exception pathways, actionable comments, consultation access, utilization dashboards, and protected staff time shared with infectious disease and antimicrobial-stewardship teams.

Limits / cautions. Voluntary self-report survey with a modest sample; responses describe perceived practice and priorities rather than measured clinical outcomes.

05 MLS curricula are adopting molecular testing faster than computational practice

Muslu et al. | Journal of Microbiology & Biology Education | May 27, 2026 | Peer-reviewed; open access | [SOURCE](#)

Finding. A review of 489 U.S. program websites found molecular-diagnostics coursework in 46.3%, but explicit bioinformatics and AI content in only 0.8% and 0.2%. Among 67 responding directors, limited curriculum time (86.6%), faculty expertise (68.7%), budget (68.7%), and infrastructure (67.2%) were leading barriers.

Why it matters. Future-ready MLS education needs observable abilities in data provenance, pipeline interpretation, algorithm validation, bias, cybersecurity, and human escalation. Modular faculty development and shared cases may be more feasible than adding entire new courses.

Limits / cautions. Course titles and public websites may undercount embedded content; the director survey response rate was about 14%; no graduate-competence or employer-outcome measures were tested. No conflicts were declared.

06 Gamification improved one form of laboratory reasoning, not all of it

Sadeh and Keshmiri | BMC Medical Education | August 19, 2026 | Peer-reviewed; open access | [SOURCE](#)

Finding. In two consecutive microbiology-internship cohorts (25 students each), a mobile gamified adjunct improved satisfaction and a puzzle-based integrative-reasoning score, but produced no added benefit on the key-feature examination.

Why it matters. Educational technology should be evaluated against multiple outcomes. MLS educators could separate engagement, pattern integration, prioritization, troubleshooting, and transfer to authentic bench decisions instead of treating a single score as proof of general clinical reasoning.

Limits / cautions. Small, single-setting, nonrandom cohort allocation; short-term assessments; no workplace-performance or retention outcome. The authors declared no competing interests.

07 Endocrine biomarkers require different expectations for serial change

Stylianou Lerpold et al. | Clinical Chemistry | August 17, 2026 | Peer-reviewed | [SOURCE](#)

Finding. Weekly sampling of 30 healthy volunteers for ten weeks showed high within-person variation for aldosterone (48.8%) and wide reference-change limits, while metanephrine varied less (13.5%). Every evaluated biomarker had an index of individuality below 1.

Why it matters. Population reference intervals alone may be weak tools for serial endocrine results. Laboratories can use biological-variation data to teach reference change values, set analytical performance goals, and help clinicians distinguish expected fluctuation from meaningful change.

Limits / cautions. Small healthy cohort under standardized conditions; results may not generalize to disease, medication exposure, older populations, or other assays. Few postmenopausal women were included.

Unexpected Connections & Leads

Established evidence is separated from the proposed collaboration or research lead.

Wastewater proficiency testing exposes a familiar laboratory-quality problem

Dutcher et al. | Emerging Infectious Diseases / CDC | August 10, 2026 | [SOURCE](#)

Established. Across two proficiency-testing rounds involving 38 laboratories, replicate respiratory-virus concentrations spanned roughly two orders of magnitude. Unit conversion, dilution, and processing-factor errors sometimes changed reported concentrations by orders of magnitude.

Collaboration lead. Clinical molecular laboratories, public-health laboratories, environmental engineers, and PT providers could adapt familiar traceability, commutability, calculation verification, and peer-group review practices to wastewater surveillance.

Next question. Which common calibrators, recovery controls, metadata fields, and reporting units would reduce between-laboratory variation without forcing one extraction method?

Boundary. Early program results with heterogeneous methods and limited same-workflow sample sizes; improved comparability has not yet been linked to better public-health decisions.

Integrated-circuit microfluidics could turn four assays into one bedside panel

Boilla et al. | Chemical Engineering Journal | September 1, 2026 | [SOURCE](#)

Established. A pump-free device separated 6 microliters of plasma from 20 microliters of whole blood and measured NT-proBNP, fibrinogen, troponin I, and CRP in about 12 minutes. In ten clinical samples, 37 of 40 biomarker-level classifications agreed with hospital assays.

Collaboration lead. Clinical chemists, coagulation specialists, nurses, biomedical engineers, and device designers could jointly define preanalytic tolerances, QC materials, reference-method comparison, operator training, and failure messages before a larger trial.

Next question. Does performance remain acceptable across hematocrit extremes, hemolysis, lipemia, anticoagulants, near-cutoff samples, reagent lots, and real bedside users?

Boundary. Proof-of-concept with only ten clinical samples; categorical agreement is not full analytical validation or evidence of better outcomes.

A veterinary sedative in the fentanyl supply creates a One Health toxicology gap

Tamama, Morrow, and Rates | Clinical Chemistry | August 17, 2026 | [SOURCE](#)

Established. High-resolution LC-qToF-MS characterized medetomidine and multiple urinary metabolites in clinical specimens. The report notes that parent drug may be absent even when metabolites are detectable, while routine urine immunoassays generally do not include medetomidine.

Collaboration lead. Clinical toxicology, emergency medicine, veterinary pharmacology, poison centers, and public-health surveillance programs could develop a metabolite-aware case definition and a shared reference-material strategy.

Next question. Which metabolite combination offers the best exposure window and how should laboratories communicate a result when clinical thresholds are not established?

Boundary. Research letter and specialized high-resolution workflow; it does not establish prevalence, a routine cutoff, or outcome benefit. Publisher access may require a subscription.

References

Every source cited in this issue is listed below with a full clickable direct link.

1. Lu S-s, Guo Y-h, Liu D-m, et al. AI-assisted digital peripheral blood morphology multicenter randomized paired method clinical validation study. npj Digital Medicine. Published August 5, 2026. doi:10.1038/s41746-026-03089-0. <https://doi.org/10.1038/s41746-026-03089-0>
 2. Jackson AN, Wyness SP, La'ulu SL, Boyd JM, Johnson-Davis KL. Increased False Positives vs Increased Interference Flags: Comparison of 2 Ethyl Glucuronide Immunoassays. The Journal of Applied Laboratory Medicine. Published August 19, 2026. doi:10.1093/jalm/jfag114. <https://doi.org/10.1093/jalm/jfag114>
 3. Jenkins TC, Young HL, Wyles DL, et al. Effects of a diagnostic stewardship intervention to de-implement widespread use of a rapid respiratory multiplex PCR test. Microbiology Spectrum. Published July 28, 2026. doi:10.1128/spectrum.00364-26. <https://doi.org/10.1128/spectrum.00364-26>
 4. She RC, Larkin P, Parrott JS, et al. Diagnostic stewardship perceptions and practices among microbiology laboratories: a national survey from the American Society for Microbiology. Journal of Clinical Microbiology. Published online June 22, 2026;64(8):e00239-26. doi:10.1128/jcm.00239-26. <https://doi.org/10.1128/jcm.00239-26>
 5. Muslu AM, Thompson C, Williams J, Basbas CG, D'Antonio EL. Advanced molecular detection, bioinformatics, and artificial intelligence era in medical laboratory science education. Journal of Microbiology & Biology Education. Published May 27, 2026. doi:10.1128/jmbe.00334-25. <https://doi.org/10.1128/jmbe.00334-25>
- Source and link verification. All hyperlinks were checked as HTTPS DOI resolvers, official journal or publisher pages, or CDC pages. URL shorteners, unknown file hosts, reposts, search-result links, and social-media claims were excluded. This materially reduces risk, although no external site can be guaranteed permanently risk-free. Access labels may change. Findings are educational intelligence, not patient-specific medical advice.

6. Sadeh M, Keshmiri F. Effect of a gamified mobile-based application on clinical reasoning skills and learning satisfaction among medical laboratory students compared with routine internship training: a quasi-experimental study. BMC Medical Education. Published August 19, 2026. doi:10.1186/s12909-026-09988-w. <https://doi.org/10.1186/s12909-026-09988-w>
7. Stylianou Lerpold A, Chowdhury R, Roys EA, et al. Biological Variation of Adrenal-Related Endocrine Biomarkers Used for Investigation of Secondary Hypertension. Clinical Chemistry. Published August 17, 2026. doi:10.1093/clinchem/hvag076. <https://doi.org/10.1093/clinchem/hvag076>
8. Dutcher HA, Antkiewicz DS, Roguet AJ, Shafer MM. Implementation and Early Outcomes of Laboratory Proficiency Testing Program for National Wastewater Surveillance System, United States, 2024. Emerging Infectious Diseases. Published August 10, 2026;32(13). doi:10.3201/eid3213.251612. <https://doi.org/10.3201/eid3213.251612>
9. Boilla SK, Lee I, Tsai WC, Tsai TH, Lee GB. Time-domain integrated-circuit biosensors for multiplexed detection of cardiovascular biomarkers via capillary microfluidics. Chemical Engineering Journal. September 1, 2026;543:178086. doi:10.1016/j.cej.2026.178086. <https://doi.org/10.1016/j.cej.2026.178086>
10. Tamama K, Morrow D, Rates T. Detection of Multiple Medetomidine Metabolites in Clinical Urine Specimens by Liquid Chromatography-Quadrupole Time-of-Flight Mass Spectrometry (LC-qToF-MS). Clinical Chemistry. Published August 17, 2026. doi:10.1093/clinchem/hvag096. <https://doi.org/10.1093/clinchem/hvag096>